

Detection of control genes

PCR results are obtained in the participating laboratories with all samples containing either DNA stock solution (DNA) or preparation from paraffin slices (P).

Lab		A		B		C		D	
Sample	DNA* (µg/ml)	DNA	P	DNA	P	DNA	P	DNA	P
1	150	+	+	-	-	+	+	(+)	+
2	30	+	+	-	+	+	+	(+)	+
3	50	+	+	+	+	+	+	+	+
4	170	+	+	+	+	+	+	+	+
5	105	+	+	+	+	+	+	+	+
6	206	+	+	+	+	+	+	+	+
7	70	+	+	+	+	+	+	+	+
8	90	+	+	-	+	+	+	-	+
9	370	+	+	+	+	+	+	+	+
10	325	+	+	+	+	+	+	+	+
11	680	+	+	+	+	+	+	+	+
12	130	+	+	+	+	+	+	+	+
13	151	+	+	+	-	+	+	+	(+)
14	181	+	+	+	+	+	+	+	+
15	480	+	+	+	+	+	+	+	+
16	422	+	+	+	+	+	+	+	+
17	104	+	+	+	+	+	+	+	+
18	107	+	+	+	+	+	+	+	+
19	447	+	+	+	+	+	-	+	+
20	131	+	+	+	+	+	+	+	+
21	92	+	+	-	-	-	+	+	(+)
22	140	+	+	+	+	+	+	+	+
23	33	+	+	+	+	+	+	+	+
24	145	+	+	+	+	+	+	+	+
25	15	+	+	+	+	+	+	(+)	+
26	120	+	+	+	+	+	+	+	+
27	10	+	+	-	+	+	+	-	(+)
28	360	+	+	+	+	+	+	+	+
29	100	+	+	+	+	+	+	+	+
30	205	+	+	+	+	+	+	+	+
31	15	+	+	+	+	+	+	+	+
32	280	+	+	+	+	+	+	+	+
33	none	n.t.	-	n.t.	+	n.t.	+	n.t.	+
34	none	n.t.	+	n.t.	+	n.t.	+	n.t.	+

Legend: DNA* concentration of DNA stock solution; A, Berlin; B, Dresden; C, Munich; D, Borstel;
+, clearly positive; (+), weakly positive; -, negative; n.t., not tested (no DNA sent to the labs);